

WAT-1 is the tool to measure drug withdrawal in pediatric patients receiving opioid and benzodiazepine sedation. 11 items are assessed on 12-point scale.

3.6 Withdrawal Assessment Tool-1 (WAT-1)

WAT-1 (;t; , ;it- t";;it--(f_J-6 -U(i: « Y ')' 7 Ht" Y(J) j=lz (i: J: '@ ft ,tTl,'@ mYc(J) filtfilt
iSJZ{ofi9 -8'Y-JL,c'9 o l JJ§ 12, , c-i!JZ{illi G *9²⁸1o

12u r.IIMti' s C7J ;:;!1' c. Ii,s C7J ffiffi	1,\ffil/7.1< ffl!	tJL,=O @-@=1
	!!!P±/tt,1? /P±	tJL,=O @-@=1
	ft;;,lff. >37 .8°C	tJL,=O -@=1
2?tr.IIC7J IliJ jji7e7.)ffil,	JUW ml	SBS O t, 'M/II!! /ti:\'b'=0 SBS +1 t,, M/1'1 = 1
	mm	tJ L,/ff!J!'.=O 9'1 /J!./ JJ!'.=1
	* ;f(9t,1"'{t,)	tJL,=O -@=1
	t) (7Jf,J:l,\Jit/J&: iit/J	tJL,/ff!J!'.=O 9'1 /J!./ IJ!'.=1
	115< 7I!< l,t,j.	1 @JtCF=O 2@Jl-:Z,t= 1
1分間の刺激観察	?''Y7'-l.:7J? < IJ9 f-	t,J: L,/ff!J!'.=O 9'1 /J!./ IJ!'.=1
	Wi	:i:E'.%=0 :!ffilJD=1
刺激後の回復	:iffif.J: (ffit->t, t);:ml (SBS 0) l: *c' C7J r.II	2分未満=0 2-5分=1 5分超=2

!ffil: Pediatr Crit Care Med 2006;7(2):107-114 fJ 1',0) fpJ>a:f c fi ;M: ffl L,iJJ
i11 C. il'iffi: !ff \$a, ftE/il=|' , r iiJJoc, lll::<tJl*, J,lxp 'fl'.# ...tffeflll
fA:&:::J' "¥' ml* f,r ,*9"n'ffil '¥5:i'!ff

WITHDRAWAL ASSESSMENT TOOL (WAT-1):J:1111

e WAT-1 (;t; , ;it- t";;it--(:t G < (;t; « Y ')' 'y'iz' l::' Ytf g(5 E3 mx.-8tcJ-c) GT;_i ,L :t G < (;t; *5cj:9'.
ntcm , 11'U(J): .mli *? tc E31J6 iSJZffli9 '8o ff(J)J:lz tJ'61E32@J72ff-¥F13iisJZ1ffii9-80

e The Withdrawal Assessment Tool (WAT-1 HiSBS⁷ c:ti: tc:l-'tcJ- < c: :t I 2B Fl3i ;::: c: tc: *J 0 i/R G iSJZlilli
9 -8« c-iff> -8 (8: 00 c: 20:00±2ff-¥F13itcJ- c) oSBS⁷iSJZloli(i:tl l, \-8WII;fH;t; Aftfil t 9-8WII;%c: GT
ff Mi:iJtm

:!=/H tJ,5' ¥&1.t A-¥9l>o (nta:fU O)il!Hic E 5 c'=t Pifmo)

- i<);01,,{Jli:/7.k:tiil'i: 121F-¥Fsmgp.c:f1:>01,,{Jli:/7.k:tiil'i:(t.fu) 0) 2, tf'l,c \,0 ;3-1J22-a :INJ\, \ ;3-0,<2-a
- O[!O±<C:/J_ /l± %; 121F-¥Fsil;:zci:log[!ll±<C:/J_ /O± %;0) 2, :IJ, :/1,c \,0 ;3-1,'2-a 2 :IJ,fJ\, \ ;3-0,'2-a
- i;:;Jli>37.8°C:121F-¥Fsmgp.c: t, < 2 :/1,n,0-(\$;:Jli:1J,37.8°C1ri'ilix.0 -8-1,<2-a-t" '5c-tJ,1 ;3-0,<2-a

27tria10)fU MO)fJ!.

- ti: J: IJ;jj(filr0)25tFsiO)til1Jll.c: :/J'1"tj(c'a60 ;3-(SBS¹:+ 1) L<2-a
Ji /l!ll!Vffl'\>:/J'jh1,f.Ji¥]"c'iP.) ;3-(SBS¹ 0)0,'2-a
- tmJil;l': *IJ;jj(ITTf0)2:nFsiO)il, "Z, q; J:l:1J,m/10)tm :IJ,11, :/1, ;3-1,<2-atJ L, /ffil*tdctFM i¥.J"Z'a60 -8-0,<2-a
- \$B:f:*IJ;jj(ITOT)f25tFsiO)il, "Z, t>cf:IJ,CB \$B:fL,C \,0;3-1,'2-a \$B:fL,C \,0 \, \ ;3-0,'2-a
- * C. * IJ)0)1, iUJJ/ & iHJ.J:mlJ; 90 ITTf0)2-5:HsiO)il, "Z, 't' II *t: .ctm/10)* C. * fJ)0)1, tLJ/ & iHJJ, @,cJ:l.ilHNO,
-¥ J,\$7J,\$7 t:O,Jl ij & IJ 9fJ:C'a:il, L,t.: ;3-1,'2-a1E <)m\, C*tdHlat;)3-0,'2-a
- a6<cf/<1, >a;:*IJ;jj(9 0 ITTf0)2-5:HsiO)il, 'c', 2@Jl-i...ta6<cf\>> <1, >a;:IJ,ffil, :/1,t.: S L<2-aO@1, * tdct1 @J0 ;3-0,'2-a

17J'ria10) flJ; fJ!,

- 'Y 'yr ,c: (-:; < 0)9 O:/ir: Y Y T ,c: (J-; < 0)9 :/ir: IJ*IJ;j&q;:c: 't' /c:, * tdct il!LJ,c:iE?, t.: ;3-1,s(a
1E< ffl\1,,(Udctffil) -8-0,<2-a
- w i s : *IJ;l&cp.c: w i s :IJ:tl!JJ.i□'90 -8-1,<2-ai:E;si;tJ -8-0,sa

fU f O)@lffi:

- ,P:ttJt<: J(S B S O),c: 0* 'Z'O)IF-fFsi: .P:f.f.J:t/<: (SBS O),c: 0* c'O)iF-fFsi: IJ, IJ; ff5:n'i!il0 ;3-2,<2-a
2-5:J'J' S 5-5:J0) -;3-1,'2-a2-5:J*:iilij0) -;3-Q,s(c

11項目で合計WAT-1スコアを取る(0-12点)

I Curley et al. State behavioral scale: A sedation assessment instrument for infants and young children supported on mechanical ventilation. Pediatr Crit Care Med 2006;7(2):107-114.

© 2007 Linda S. Franck and Martha A.Q. Curley
Translated by: Hisae Kaneko, January 24, 2019



Made available under a Creative Commons Attribution-NonCommercial-NoDerivs License (CC BY-ND 2.0)
Full terms of license available here: <https://creativecommons.org/licenses/by-nd/2.0/>