

WAT-1 is the tool to measure drug withdrawal in pediatric patients receiving opioid and benzodiazepine sedation. 11 items are assessed on 12-point scale.

3.6 Withdrawal Assessment Tool-1 (WAT-1)

WAT-1 (;t; , ;it- t";;it--(fJ_6 -U(i: « Y ')')'yiz' l::°Ytf g(5 E3 mx.-8tcJ-c) GT;_i ,L :t G < (;t; *5cj:9'.
 iSJZ{ofi9 -8'Y-JL,c'9 o l l JJ§ 12, , c-i!JZ{illi G *9²⁸1o

12u r.IIMti' s C7J ;:;!1' c. Ii,s C7J ffiffi	1,\ffil/7.L< ffl!	tJL,=O @-@=1
	!!!P±/tt,1? /P±	tJL,=O @-@=1
	ft;;,lff. >37 .8°C	tJL,=O -@=1
2?tr.lIC7J lliJ jji7.)ffil,	JUW ml	SBS O t, 'M/II!! /ti:\'b'=0 SBS +1 t,, M/1'1 = 1
	mm	tJ L,/ff!J!'.=O 9'1 /J!./ JJ!'.=1
	* ;f(9t,1"'t,.)	tJL,=O -@=1
	t) (7Jf,J:l,Jit/J&: iit/J	tJL,/ff!J!'.=O 9'1 /J!./ IJ!'.=1
	115< 7l!< l,t,j.	1 @JtCF=O 2@Jl.:Z,t= 1
1分間の刺激観察	?''Y7'-l.:7J? < IJ9 f-	tJ: L,/ff!J!'.=O 9'1 /J!./ IJ!'.=1
	Wi	:i'E.%=0 :ffilJD=1
刺激後の回復	:iffif.J: (ffit->t, t);:ml (SBS 0) l: *c' C7J r.II	2分未満=0 2-5分=1 5分超=2

1,ffil:Pediatr Crit Care Med 2006;7(2):107-114 fJ 1',0) fpJ>a:f c fi ;M: ffl L,iJJ
 ill C. il'iffi: !ff \$a, ftE/il=ll' , r iiJoc,II!::<tJl*, J,lxp 'fl'.# ...tffefllll
 fA;&:::J' "¥' ml* f,r ,*9"n'ffil '¥5:i'!ff

WITHDRAWAL ASSESSMENT TOOL (WAT-1):J:1111

- e WAT-1 (;t; , ;it- t";;it--(:t G < (;t; « Y ')')'yiz' l::°Ytf g(5 E3 mx.-8tcJ-c) GT;_i ,L :t G < (;t; *5cj:9'.
 ntcml , 11'U(J): .mli *?tc E31J6 iSJZffil9 '8o ff(J)J:lz tJ'61E32@J72ff-¥F13iisJZ1ffii9-80
- e The Withdrawal Assessment Tool (WAT-1 HiSBS⁷ c:ti: tc:l-'tcJ- < c: :t I 2B Fl3i ;::: c: tc: *J 0 i/R G iSJZilli
 9 -8« c-iff> -8 (8: 00 c: 20:00±2ff-¥F13itcJ- c) oSBS⁷iSJZ1oli(i:tl l, \-8WII;fH;t; Aftfil t 9-8WII;%c: GT
 ff Mi:iJtm

:!=/H tJ,5' ¥&1.t A-¥9l>o (nta:fU O)il!Hic E 5 c'=t Pifmo)

- i<);01, {Jli:7.k:tiil'i: 121F-¥Fsmg.c:f1:>01, {Jli:7.k:tiil'i:(n.fu) 0) 2, tf'l,c \,0 ;3-1J22-a :INJ\, \;3-0,<2-a
- O[!O±<C:J_3 /l± %; 121F-¥Fsil; zci: log[!ll±<C:J_3 /O± %;0) 2, :IJ, :/1,c \,0 ;3-1,'2-a 2 :IJ,fJ\, \;3-0,'2-a
- i;,\$;Jli>37.8°C:121F-¥Fsmg.c: t, < 2 :/1,n,0-(\$;Jli:1J,37.8°C1ri'ilix.0 -8-1,<2-a-t' '5c-tJ,1 ;3-0,<2-a

27tria10) fU MO)fJ!.

- ti: J: IJ;jj(filr0)25tFsiO)til1Jll.c: :/J'1"tj(c'a60 ;3-(SBS¹:+1) L<2-a
 Ji /ll!!!Vffl'\>:/jhl,f.Ji¥]"c'iP.) ;3-(SBS¹ 0)0,'2-a
- tmJil;l': *IJ;jj(ITTf0)2:nFsiO)il, "Z, q; J:l:1J,m/10)tm :IJ,11, :/1, ;3-1,<2-atJ L, /ffil*tdctFM i¥.J"Z'a60 -8-0,<2-a
- \$B:f:*IJ;jj(ITOT)f25tFsiO)il, "Z, t>cf:IJ,CB \$B:fL,C \,0;3-1,'2-a \$B:fL,C \,0 \, \;3-0,'2-a
- * C.* IJ)0)1, iUJJ/ & iHJ.J:mlJ; 90 ITTf0)2-5:HsiO)il, "Z, 't' II* t: .ctm/10)* C.* fJ)0)1,, tJ/ & iHJJ, @,cJ:l.ilHNO,
 -¥ J,\$7J,\$7 t:O,Jl ij & IJ 9fJ:C'a:il, L,t.: ;3-1,'2-a1E <)m\ C*tdHlat;)3-0,'2-a
- a6 < cf/<1, >a;:*IJ;jj(9 0 ITTf0)2-5:HsiO)il, 'c', 2@Jl-i...ta6 < cf\>> <1, >a;:IJ,ffil, :/1,t.: S L<2-aO@1, * tdct1 @J0 ;3-0,'2-a

17J'ria10) flJ; fJ!,

- 'Y 'yr ,c:(j-; <)9 O:/ir:Y YT ,c:(j-; <)9 /ir:IJ*IJj&q;:c:'t' /c:, * tdct il!LJ,c:iE?, t.: ;3-1,s(a
 1E< ffl\1,(Udctffil) -8-0,<2-a
- w: s :*IJ;l&cp,c:w: s :IJ;t!JJ.i□'90 -8-1,<2-ai:E;si;tJ -8-0,sa

fU f O)@lffi:

- ,P:ttJt<: J(S B S O),c: 0*'Z'O)IF-fFsi: .P:f.f.J:t/<: (SBS'O),c: 0*c'O)iF-fFsi:1J, IJ; ff5:n'i!il0 ;3-2,<2-a
 2-5:J'J' S 5-5:J0) -;3-1,'2-a2-5:J*:iilij0) -;3-Q,s(c

11項目で合計WAT-1スコアを取る(0-12点)

I Curley et al. State behavioral scale: A sedation assessment instrument for infants and young children supported on mechanical ventilation. Pediatr Crit Care Med 2006;7(2):107-114.

© 2007 Linda S. Franck and Martha A.Q. Curley
 Translated by: Hisae Kaneko, January 24, 2019



Made available under a Creative Commons Attribution-No Derivatives License (CC BY-ND 2.0)
 Full terms of license available here: <https://creativecommons.org/licenses/by-nd/2.0/>